

ORIGINAL
ARTICLE

Erdal Uysal¹
Efe Sezgin²
Hasan Bakir¹
Ahmet Gurur³
Mehmet Sokucu⁴

¹Sanko University School of
Medicine, Department of
General Surgery, Gaziantep,
Turkey

²İzmir Institute of
Technology, İzmir, Turkey

³Sanko Hospital,
Department of General
Surgery, Gaziantep, Turkey

⁴Sanko University School of
Medicine, Department of
Pathology, Gaziantep,
Turkey

Corresponding Author:

Erdal Uysal
Sanko University School of
Medicine, Department of
General Surgery, Gaziantep,
Turkey
Tel: +90 342 211 50 00
dredaluyisal@hotmail.com

Received: 03.07.2017
Acceptance: 14.09.2017
DOI: 10.18521/kt.325651

Konuralp Medical Journal
e-ISSN1309-3878
konuralptipdergi@duzce.edu.tr
konuralptipdergisi@gmail.com
www.konuralptipdergi.duzce.edu.tr

Is Routine Histopathological Gallbladder Examination Necessary After Cholecystectomy? Evaluation of the Results of 1366 Cholecystectomy Specimens in Single Center

ABSTRACT

Objective: It was aimed to evaluate the results of routine histopathological examination after cholecystectomy and to investigate the necessity of routine histopathologic examination after cholecystectomy.

Methods: The study was designed retrospectively. One thousand three hundred sixty six patients who underwent laparoscopic and open cholecystectomy at Private Sani Konukoglu Hospital with pre-diagnosis of benign gallbladder disease between November 2011 and May 2017 were included in the study. Patients' demographic data, pathologic results, macroscopic appearance of the specimen, and cancer staging were recorded. The distribution and frequency of pathologic diagnoses and the prevalence of incidental gallbladder cancer (GBC) were evaluated. Pathologic findings were compared in terms of age groups and gender relations.

Results: The number of patients included in the study was 1366. Diagnosed with chronic cholecystitis patients were 1,303 (95%), 39 (3%) with acute cholecystitis, 7 (0.5%) with gallbladder cancer, and 17 (1.5%) with other diagnoses of the patients. Statistical significance was found between the groups in terms of the mean age ($p = 0.0002$). Comparisons between groups in terms of cholesterolysis were statistically significant ($p = 0.0003$). There was a significant relationship between mucosa atrophy and gender ($p = 0.001$).

Conclusions: The histopathological spectrum of gallbladder is quite extensive. Incidental GBC may not be detected by preoperative imaging methods. Incidental GBC are usually asymptomatic. T₂, T₃ and T₄ GBC were also encountered in our study. All of these patients need additional operations. In the absence of routine histopathologic examination, metastatic advanced GBC may be encountered because no treatment plans could make. Thus, we do recommend routine histopathological examination.

Keywords: Cancer, Cholesterolysis, Dysplasia, Gallbladder, Histopathologic, Metaplasia.

Kolesistektomi Sonrası Rutin Histopatolojik Safra Kesesi İncelemesi Gerekli Mi? Tek Merkezdeki 1366 Kolesistektomi Spesmeninin Sonuçlarının Değerlendirilmesi

ÖZET

Amaç: Kolesistektomi sonrası rutin olarak yapılan histopatolojik inceleme sonuçlarının değerlendirilmesi, kolesistektomiler sonrası rutin histopatolojik incelemenin gerekliliğini araştırmak amaçlanmıştır.

Gereç ve Yöntem: Çalışma retrospektif olarak dizayn edilmiştir. Bening safra kesesi hastalığı ön tanısı ile Kasım 2011- Mayıs 2017 yılları arasında Özel Sani Konukoğlu Hastanesinde laparoskopik ve açık kolesistektomi uygulanan 1366 hasta çalışmaya dahil edildi. Hastaların demografik verileri, patoloji sonuçları, spesmenin makroskopik görünümü, kanser evrelemesi kayıt edildi. Patolojik tanıların dağılımı ve sıklığı, incidental safra kesesi kanseri prevalansı değerlendirildi.

Bulgular: Çalışmaya dahil edilen hasta sayısı 1366 idi. Hastaların 1303 (%95)'ü kronik kolesistit, 39 (%3)'ü akut kolesistit, 7 (%0.5)'i safra kesesi kanseri, 17 (%1.5) hastada diğer tanılar tespit edildi. Gruplar arasında yaş ortalaması açısından istatistiksel anlamlılık saptandı ($p=0.0002$). Kolesterolizis açısından gruplar arasında yapılan karşılaştırmalarda istatistiksel olarak anlamlılık bulundu ($p=0.0003$). Mukoza atrofisi ile cinsiyet arasında anlamlı ilişki tespit edildi ($p=0.001$).

Sonuç: Safra kesesi spesmenlerinin histopatolojik incelenmesinde en sık görülen tanı kronik kolesistittir. Ancak kolesistektomi sonrası, safra kesesinin histopatolojik spekturumu oldukça geniştir. Incidental safra kesesi tümörleri preoperative görüntüleme yöntemleri ile tespit edilemeyebilir. Incidental safra kesesi tümörleri genellikle asemptomatik seyretmektedir. Çalışmamızda T₂, T₃ ve T₄ safra kesesi tümörlerine de rastlanılmıştır. Bu hastaların tümüne ek girişimler gerekmiştir. Rutin histopatolojik inceleme yokluğunda, tedavi planı yapılamadığından metastatic ileri evre safra kesesi tümörleriyle karşılaşılabilir. Bu nedenle rutin histopatolojik incelemenin yapılmasını önermekteyiz.

Anahtar Kelimeler: Safra kesesi, Histopatoloji, Kanser, Kolesterolizis, Displazi, Metaplazi

INTRODUCTION

Cholecystectomy is performed electively or urgently in gallbladder diseases. The most important aspect in the histopathological examination of the gallbladder specimens is the malign tumors of the gallbladder. Although the malignant diseases of the gallbladder are rarely seen, they are quite aggressive. In one study, incidence in western societies was reported as 0.9 and 1.9 in 100,000 men and women, respectively (1). However, it is observed in northern India and Pakistan with a high incidence of 21.5 and 13.8 in 100,000 cases, respectively (2, 3).

Since there are no specific signs and symptoms of gallbladder cancer (GBC), they usually appear late stage of the cancer. Therefore, the diagnosis can usually be made in the advanced stage. While 5-year survival is below 5%, overall mean survival is 6 months (4).

Cholecystectomy is sufficient for curative therapy in the mucosa limited (Tis) and T1a GBCs (5, 6). Early stage GBCs are usually incidentally detected during histopathological evaluation of the pathologist's cholecystectomy specimen. GBC diagnosis are made as incidental in the histopathological examination of the gallbladder specimen which is thought to be benign in 60-80% of cases. Incidental primary GBC ratio was reported as 0.19-3.3% (7, 8). It has been reported that the prognosis is better in incidental GBC. It was shown that this is because of early diagnosis and fast definitive treatment (9, 10).

Histopathologic examination is very important when it comes to the confirmation of the clinical and radiological diagnosis. Histopathologic examination is also an important prognostic tool in advanced treatment of patients. Histopathological examination also plays an important role in medicolegal cases. Although histopathological examinations are routinely performed in many centers, they are performed selectively in some centers (9, 11). Routine histopathological examination of the gallbladder is still a controversial subject. In some publications, it was emphasized the importance of the routine histopathological examination in cancer diagnosis, whereas in some publications, selective histopathological examination is recommended (12, 13). It has been reported that in patients with macroscopically abnormal gallbladder, elderly, female, high-risk group with high alkaline phosphatase levels; selective histopathologic examinations have higher cost effective. In addition, selective histopathological examinations are thought to reduce the workload of pathologists (14). The College of American and the UK Royal College of Pathologists recommend selective histopathologic examination of surgical specimens (15).

In this study, it was aimed to evaluate the results of routine histopathological examination

after cholecystectomy and to investigate the necessity of routine histopathologic examination after cholecystectomy.

MATERIAL AND METHODS

The study was designed retrospectively. One thousand three hundred sixty six patients who underwent laparoscopic and open cholecystectomy at Private Sani Konukoglu Hospital with pre-diagnosis of benign gallbladder disease between November 2011 and May 2017 were included in the study. Patients with imaging findings consistent with preoperative gallbladder cancer were not included in the study. Cholecystectomies made in addition to liver resection or pancreaticoduodenectomy were not included in the study as well. All gallbladder specimens were sent to the pathology department. Patient data were retrospectively obtained by recording the data in patients' files and electronic records based on the protocol created initially for this study. Patients' demographic data, pathologic results, macroscopic appearance of the specimen, and cancer staging were recorded. The distribution and frequency of pathologic diagnoses and the prevalence of incidental GBC were evaluated. Pathologic findings were compared in terms of age groups and gender relations. Staging of gallbladder tumors was performed according to the American Joint Committee on Cancer TNM system. In this research which is not invasive approval from the ethical committee was not requested. Retrospective file scanning was performed. The study was conducted in accordance with the Declaration of Helsinki. This study is registered at researchregistry.com by the number ID researchregistry 3114.

Statistical Analysis: Distribution of categorical factors (such as metaplasia and cholesterolosis groups) among the patients were compared by Chi-square tests. Age was analysed as continuous variable with a normal distribution and expressed as mean \pm standard deviation. ANOVA was used to test the age difference between cholecystitis and other groups. P-values less than 0.05 were considered as statistically significant. All statistical analyses were conducted by SAS/STAT version 9.3 (SAS Institute, Inc, Cary, North Carolina, USA).

RESULTS

The number of patients included in the study was 1,366. Diagnosed with chronic cholecystitis patients were 1,303 (95%), 39 (3%) with acute cholecystitis, 7 (0.5%) with gallbladder cancer, and 17 (1.5%) with other diagnoses of the patients. Female and male patients were 954 (69.8%) and 412 (30.1%) respectively (Table 1). Statistical significance was found in gender comparison ($p = 0.0001$). The mean age in the chronic cholecystitis group was 49 ± 15 , 59 ± 14 in the acute cholecystitis group and 52 ± 18 in the other pathologic diagnosis group (Figure 1-2).

Table 1. Diagnosis and age groups distribution

Diagnosis	N (%)
Chronic cholecystitis	1303 (95%)
Acut cholecystitis	39 (3%)
Eosinophilic cholecystitis	4 (0.3%)
Gallbladder Cancer	7 (0.5%)
polyp	12 (0.9%)
Xanthogranulomatous cholecystitis	1 (0.1%)
Age Groups	
<25	67 (5%)
26 – 45	477 (35%)
46 – 50	153 (11%)
> 50	669 (49%)

Statistical significance was found between the groups in terms of the mean age ($p = 0.0002$). Histopathological examination of gallbladder revealed metaplasia in 16 (1.1%) patients. There was no statistical significance seen between the groups in terms of metaplasia ($p = 0.92$). There was no statistical significance between the metaplasia and age groups and gender ($p = 0.99$) ($p = 0.81$). Dysplasia was observed in 3 patients (0.2%) in the chronic cholecystitis group. Diffuse cholesterolysis was detected in 365 (26.7%) and cholesterol polyp in 28 (2%) of the examined gallbladder. Comparisons between groups in terms of cholesterolysis were statistically significant ($p = 0.0003$). Comparisons between cholesterolysis and age groups and gender were statistically significant ($p = 0.0001$) ($p < 0.0001$). Mucosal atrophy was

observed in 208 (15%) of the examined gallbladder (Table-2). There was no statistically significant difference between the groups in terms of mucosa atrophy ($p = 0.32$). However, there was a significant relationship between mucosa atrophy and gender ($p = 0.001$) (Table 3).




Figure 1. Distribution of gallbladder pathology among the age groups.




Figure 2- Distribution of male and female patients among the age groups.

Table 2. Relationship between diagnosis and metaplasia, dysplasia, cholesterolosis, mucosa atrophy

	Diagnosis			P
	Chronic cholecystitis (n=1303)	Acut cholecystitis (n=39)	Other (n=24)	
Age (mean±SD)	49±15	59±14	52±18	0.0002
Sex (%)				0.0001
Male (n=412)	94	5	0.5	
Female (n=954)	96	2	2	
Metaplasia				0.92
• Yes (n=16) (%)	100	0	0	
• No (n=1346) (%)	95	3	2	
Dysplasia (n=3) (%)	100	0	0	
Cholesterolosis				0.0003
• Yes (n=365) (%)	99	0.6	0	
• No (n=967) (%)	94	4	2	
• Cholesterol polyp (n=28) (%)	100	0	0	
Mucosa atrophy				0.32
• Yes (n=208) (%)	97	3	0.5	
• No (n=1156) (%)	95	3	2	

Table 3. Relationship between metaplasia, dysplasia, cholesterolosis, mucosa atrophy and sex, age groups.

	Metaplasia		Dysplasia + Metaplasia (n=3) %	P	Cholesterolosis		Cholesterol polyp (n=28) %	p	Mucosa atrophy		p
	Yes (n=16) %	No (n=1346) %			Yes (n=365) %	No (n=967) %			Yes (n=208) %	No (n=1156) %	
Sex				0.81				0.0001			0.001
Male	37	30	33		18	35	29		40	28	
Female	63	60	67		82	65	71		60	72	
Age Groups				0.99				<0.0001			0.16
<20	0	1	0		1	1	3		1	1	
20-30	6	9	0		12	7	11		5	9	
31-40	19	19	33		23	17	18		16	19	
41-50	19	22	33		23	22	32		26	22	
51-60	25	23	0		24	22	36		27	22	
61-70	19	16	33		3	13	0		16	17	
>70	12	9	0		4	11	0		8	9	

DISCUSSION

Routine histopathological examination of all gallbladder specimens is still a controversial issue (9, 15). Selective histopathologic examination is recommended for some publications (16). In selective histopathological examination, histopathologic examination is performed especially in case of macroscopic pathologies occurred in the perioperative gallbladder and > 60 years of age (6). Dix *et al.*, Darmas *et al.*, Ramraje SN *et al.* have proposed a selective approach. Thus, it has been argued that the workload of the pathologist will reduce and the cost will go down as well (4, 6, 15). In a study conducted in India, 47% of gallbladder specimens contained macroscopic abnormalities. It has been reported that 13 of these patients have GBC. In this study, histopathologic examination of gallbladder specimens containing only macroscopically abnormalities was proposed (14). The frequency of the GBC's in Asian countries and North India is quite high. Selective histopathologic examination in areas with high susceptibility has not been found appropriate by some researchers (17). However, Chin KF *et al.* stated that demographic characteristics did not affect selective treatment (18).

Although there are some authors who recommend selective histopathological examination, there are also authors who do not recommend it. Siddiqui *et al.* reported in their study that the histopathological spectrum of gallbladder was quite variable. It is stated that the incidence of incidental GBC is not uncommon, and subclinical malignancies cannot be detected as a result of non-routinely histopathologic examinations and this situation will be catastrophic. Thus, Siddiqui *et al.* suggests routine histopathological examination of all gallbladder specimens (11). Agarwal *et al.*

reported that late recurrences due to gallbladder specimens without histopathological examination may be seen and may cause poor prognosis due to unresectability. Therefore, it is recommended that all cholecystectomy specimens be sent to pathologic examination for histopathological examination (9). Our center has a policy to send all gallbladder specimens to a histopathological examination. However, macroscopic abnormalities of the mucosa were detected in all the specimens except for a specimen, which had 7 GBC diagnoses in our series. Moreover, the average age of the patients was 63.

Gallbladder diseases are more common in women than in men (19). In our study, 69.8% of the patients were female. Our findings were compatible with the literature.

Chronic cholecystitis was found to be the most common diagnosis in our study. Siddiqui FG *et al.* (11) reported that the diagnosis of chronic cholecystitis was 92.3% and Basak F *et al.* (12) stated that it was 96.3%. In our study, the diagnosis of chronic cholecystitis was found to be 95% in agreement with the literature.

Histopathological examination of gallbladder revealed a mucocoeles incidence of approximately 3% (11). No mucocoeles were found among the specimens examined in our study. Xanthogranulomatous cholecystitis is an inflammatory disease characterized by the accumulation of lipid-laden macrophages, fibrous tissue, acute and chronic inflammatory cells. It may mimic gallbladder cancer as involvement in neighboring organs (20). It has also been reported that there is a relationship between xanthogranulomatous cholecystitis and carcinoma of the gallbladder cancer (21). In our study,

xanthogranulomatous cholecystitis was detected in a specimen. Eosinophilic cholecystitis is a rare form of Cholecystitis. Histopathologically, it is characterized by transmural leukocyte infiltration with more than 90% eosinophil. Although etiology is uncertain, it has been reported that it can be associated with allergies, parasites, hypereosinophilic syndromes, eosinophilic gastroenteritis, cholelithiasis, and acalculous cholecystitis (22). In our study, eosinophilic cholecystitis was detected in 3 specimens.

The incidence of gallbladder polyps varies between 4.6-12% (23, 24). Gallbladder polyps are often asymptomatic and are usually detected incidentally during ultrasonography. Non-neoplastic or neoplastic polyps can be seen in the gallbladder. Neoplastic polyps can be seen as adenomas, adenocarcinomas and squamous cell carcinomas (25). The most common polyps are cholesterol polyps as the rate of 60-90% (26).

In our study, the cholesterol polyp ratio was 70%. Neoplastic polyps are seen rarely. Adenomas and carcinomas are involved in neoplastic polyps. Although adenomas are usually benign, they exhibit premalignant behavior at 4% (23). In our study, adenomatous polyp was detected in 10 specimens. One of the specimens was reported as a tubulovillous adenomatous polyp. The other two specimens were reported as fibromas, while the other one was adenomyomas. American Gastrointestinal and Endoscopic Surgeons recommends the follow-up of asymptomatic polyps <5 mm, and treatment of larger single and symptomatic polyps with cholecystectomy (27). The approach in our center is in the same direction as laparoscopic cholecystectomy was suggested with gallbladder polyps > 1 cm and symptomatic disease.

Although cholesterosis of the gallbladder is common, it is a rarely reported subject. Cholecystectomy is seen in about 20% of the specimens (28). Cholesterosis rate was reported as 25% by Ozgur T *et al.* (29). In our study, Cholesterosis of the gallbladder was seen in 26%. Gallbladder cholesterosis can be correlated with elevated serum cholesterol. In addition, cholesterol that accumulates in the gallbladder wall causes gallbladder dysfunction. Cholesterosis can also cause idiopathic pancreatitis (28, 30, 31). Yaylak F *et al.* reported that cholesterosis of the gallbladder might be associated with metaplasia at the gallbladder (32). Roa I *et al.* reported that cholesterosis was significantly lower in specimens with gallbladder cancer (33). In our study, cholesterosis was not seen in specimens with gallbladder cancer. Studies conducted by Dairi S *et al.* and Yaylak F *et al.* showed that cholesterosis was significantly higher in women (28, 32). In our study, histopathological examination of gallbladder specimens revealed that cholesterosis was significantly higher in females

compared to males. This may be due to the difference in cholesterol metabolism between male and female (34). The clinical significance of cholesterosis of the gallbladder is still unclear (32). Therefore, wider series of studies are needed.

Changes caused by lymphocytes and plasma cells during inflammatory processes that occur in the mucous membranes of chronic cholecystitis result in mucosal atrophy. Mucosa atrophy was detected in 15% of the specimens in our study. Hassan *et al.* found that there was no significant association between mucosal atrophy and Helicobacter pylori in their study conducted on helicobacter pylori and gallbladder mucosal changes (35).

GBC is a very aggressive malignant disease although that is rarely seen. It especially affects the elderly population. It has a poor prognosis in the absence of early diagnosis and treatment. GBC is more common in women relative to men. It shows peak incidence in seventh decade of life (8). Approximately 0.3-2% of patients with benign cause of cholecystectomy have GBC (36, 37). This rate was 0.5% in our study. Basak F *et al.* found this rate as 0.23%, whereas the rate in Asian countries and North India is higher than in western countries (12, 38). The most important reason for this depends on variables of the GBC incidence based on the geographical region.

There are two important sequences in gallbladder carcinogenesis. One of them is metaplasia-dysplasia and carcinoma development and the other one is adenoma-carcinoma development. Metaplastic changes cause precancerous changes in gallbladder epithelium (39). The relationship between gallbladder cancer and metaplasia-dysplasia has been shown in studies (8). Basak F *et al.* showed that metaplasia is associated with carcinoma in 2 of 4 patients whose gallbladder cancer was detected in their study (12). In our study, cancer and metaplasia associations were not seen in the reports. In our study, 16 (1.1%) metaplasias were found in histopathological examinations of gallbladder. Dysplasia was observed in only 3 (0.2%) specimens.

It has long been believed that there is a relationship between GBC and chronic inflammation, which is the cause of gallbladder stones (40). Gallstones are present in 54-97% of GBCs (12). It has been reported that gallstones may cause long-term chronic inflammation of the gallbladder (41). The association between chronic inflammations, gallstones and GBC is still controversial. In our study, the mass was accompanied by gallstones in 4 (57%) gallbladder diagnosed with GBC. Our findings were consistent with the literature. GBC and acute cholecystitis have also been reported (42). However, acute cholecystitis and GBC association did not observed in our study.

In their meta-analysis, Swank *et al.* reported that tumors of the Tis or T₁ stage were the rate of 42.4% (43). Deng *et al.* have shown that 61.4% of gallbladder tumors are in T_{1a} or T_{1b} (42). Basak F *et al.* reported T₂, T₃, and T₄ gallbladder tumors as 26.4%, 24%, and 7.2%, respectively (12). In our study, Tis gallbladder tumor was found in two patients (28%), T₁ in one patient (14%), T₂ in two patient (28%), T₃ in one patient (14%) and T₄ in one patient (14%). Cholecystectomy is curative in Tis and T_{1a} biliary cancer, which is limited in the mucosa (5, 6). Five-year survival in these tumors was reported as 100% (44). Cholecystectomy by itself is not sufficient in T₂, T₃ and T₄ tumors. New surgical interventions and adjuvant treatments involving hepatic resection and lymph node dissection targeting the tumor-free surgical margin should be planned (12, 45).

There are some limitations in our study. Although our study did not include a large series, it was difficult to reach some data due to the fact that it is retrospective and some patients taking additional surgical procedures and adjuvant

treatment at other centers. These findings are from a single region of Turkey. The results in regions with high incidence of gallbladder cancer may differ. Therefore, there is a need for multicenter studies from different regions and countries.

CONCLUSION

The most common diagnosis in the histopathological examination of gallbladder specimens is chronic cholecystitis. However, after cholecystectomy, the histopathological spectrum of gallbladder is quite extensive. Incidental gallbladder tumors may not be detected by preoperative imaging methods. Incidental gallbladder tumors are usually asymptomatic. T₂, T₃ and T₄ gallbladder tumors were also encountered in our study. All of these patients need additional operations. In the absence of routine histopathologic examination, metastatic advanced gallbladder tumors may be encountered because no treatment plans could make. Thus, we do recommend routine histopathological examination.

REFERENCES

1. Cancer Statistics Registrations, England (Series MB1), No. 43, 2012. Office for National Statistics. <http://www.ons.gov.uk/ons/rel/vsob1/cancer-statisticsregistrations-england-series-mb1-/no-43-2012/> (cited June 2015).
2. Lobo L, Prasad K, Satoskar RR. Carcinoma of the gall bladder: a prospective study in a tertiary hospital of Bombay, India. *J Clin Diagn Res* 2012; 6(4):692–695.
3. Randi G, Franceschi S, La Vecchia C. Gallbladder cancer worldwide: geographical distribution and risk factors. *Int J Cancer* 2006; 118: 1,591–1,602.
4. Darmas B, Mahmud S, Abbas A, et al. Is there justification for the routine histological examination of straightforward cholecystectomy specimen? *Ann R Coll Surg Engl* 2007; 89: 238–241.
5. M.T. Hueman, C.M. Vollmer, T.M. Pawlik. Evolving treatment strategies for gallbladder cancer. *Ann Surg Oncol*. 2009;16(8):2101-15.
6. Taylor HW, Huang JK. 'Routine' pathological examination of the gallbladder is a futile exercise. *Br J Surg* 1998; 85: 208.
7. Jetley S, Rana S, Khan S et al. Incidental gall bladder carcinoma in laparoscopic cholecystectomy: a report of 6 cases and a review of the literature. *J Clin Diagn Res* 2013; 7: 85–88.
8. Kalita D, Pant L, Singh S et al. Impact of routine histopathological examination of gall bladder specimens on early detection of malignancy – a study of 4,115 cholecystectomy specimens. *Asian Pac J Cancer Prev* 2013; 14: 3,315–3,318.
9. Agarwal AK, Kalayarsan R, Singh S, et al. All cholecystectomy specimens must be sent for histopathology to detect inapparent gallbladder cancer. *HPB (Oxford)*. 2012;14(4):269-73.
10. D. Fuks, J.M. Regimbeau, Y.P. Le Treut, et al. Incidental gallbladder cancer by the AFC-GBC-2009 Study Group. *World J Surg*. 2011;35(8):1887-97.
11. Siddiqui FG, Memon AA, Abro AH, et al. Routine histopathology of gallbladder after elective cholecystectomy for gallstones: waste of resources or a justified act? *BMC Surg*. 2013;13:26.
12. Basak F, Hasbahceci M, Canbak T, et al. Incidental findings during routine pathological evaluation of gallbladder specimens: review of 1,747 elective laparoscopic cholecystectomy cases. *Ann R Coll Surg Engl*. 2016;98(4):280-3.
13. Emmett CD, Barrett P, Gilliam AD, et al. Routine versus selective histological examination after cholecystectomy to exclude incidental gallbladder carcinoma. *Ann R Coll Surg Engl*. 2015;97(7):526-9.
14. Mittal R, Jesudason MR, Nayak S. Selective histopathology in cholecystectomy for gallstone disease. *Indian J Gastroenterol* 2010; 29: 26–30.
15. Ramraje SN, Pawar VI. Routine histopathologic examination of two common surgical specimens-appendix and gallbladder: is it a waste of expertise and hospital resources? *Indian J Surg*. 2014;76(2):127-30.
16. Lohsiriwat V, Vongjirad A, Lohsiriwat D. Value of routine histopathologic examination of three common surgical specimens: appendix, gallbladder, and hemorrhoid. *World J Surg* 2009; 33: 2,189–2,193.

17. Pai SA, Bhat MG. Selective histopathology of gall bladders is unscientific and dangerous. *Surgeon* 2004; 2: 241.
18. Chin KF, Mohammad AA, Khoo YY, et al. The impact of routine histopathological examination on cholecystectomy specimens from an Asian demographic. *Ann R Coll Surg Engl.* 2012;94(3):165-9.
19. Channa NA, Soomro AM, Ghangro AB: Cholecystectomy is becoming an increasingly common operation in Hyderabad and adjoining areas. *Rawal Med J* 2007; 32(2):128–130.
20. Park JW, Kim KH, Kim SJ, et al. Xanthogranulomatous cholecystitis: Is an initial laparoscopic approach feasible? *Surg Endosc.* 2017 Jun 7. doi: 10.1007/s00464-017-5604-z.
21. Rao RV, Kumar A, Sikora SS, et al. Xanthogranulomatous cholecystitis: differentiation from associated gallbladder carcinoma. *Trop Gastroenterol* 2005;26:31–3.
22. Sánchez-Pobre PI, López-Ríos Moreno F, Colina F, et al. Eosinophilic cholecystitis: an infrequent cause of cholecystectomy. *Gastroenterol Hepatol.* 1997;20(1):21-3.
23. Matos AS, Baptista HN, Pinheiro C, et al. Gallbladder polyps: how should they be treated and when? *Rev Assoc Med Bras* 2010; 56(3):318–321.
24. Cha BH, Hwang JH, Lee SH, et al. Pre-operative factors that can predict neoplastic polypoid lesions of the gallbladder. *World J Gastroenterol* 2011; 17: 2216-22.
25. Park KW, Kim SH, Choi SH, Lee WJ. Differentiation of nonneoplastic and neoplastic gallbladder polyps 1 cm or bigger with multidetector row computed tomography. *J Comput Assist Tomogr* 2010; 34: 135-9.
26. Andrén-Sandberg A. Diagnosis and Management of Gallbladder Polyps. *N Am J Med Sci* 2012; 4: 203-11.
27. SAGES Guideline Committee. SAGES guidelines for the clinical application of laparoscopic tract surgery, section H – gallbladder polyps. January 2010. Available at <http://www.sages.org/publications/id/06/>.
28. Dairi S, Demeusy A, Sill AM, et al. Implications of gallbladder cholesterosis and cholesterol polyps? *J Surg Res.* 2016 ;200 (2):467-72.
29. T. Ozgur, S. Toprak, A. Koyuncuer, M. et al. Do histopathological findings improve by increasing the sample size in chole-cystectomies. *World J Surg Oncol.* 2013;11:245.
30. Parrilla Paricio P, Garcia Olmo D, Pellicer Franco E, et al. Gallbladder cholesterosis: an aetiological factor in acute pancreatitis of uncertain origin. *Br J Surg* 1990; 77:735.
31. Khairy GA, Guraya SY, Murshid KR. Cholesterosis. Incidence, correlation with serum cholesterol level and the role of laparoscopic cholecystectomy. *Saudi Med J* 2004;25:1226.
32. Yaylak F, Deger A, Ucar BI, et al. Cholesterosis in routine histopathological examination after cholecystectomy: what should a surgeon behold in the reports? *Int J Surg.* 2014;12(11):1187-91
33. Roa I, de Aretxabala X, Ibacache G, et al. Association between cholesterosis and gallbladder cancer. *Rev Med Chil.* 2010;138(7):804-8.
34. Wang X, Magkos F, Mittendorfer B. Sex differences in lipid and lipoprotein metabolism: it's not just about sex hormones. *J Clin Endocrinol Metab.* 2011;96(4):885-93.
35. Hassan EH, Gerges SS, El-Atrebi KA, et al. The role of H. pylori infection in gall bladder cancer: clinicopathological study. *Tumour Biol.* 2015;36(9):7093-8.
36. Dix FP, Bruce IA, Krypczyk A, et al. A selective approach to histopathology of the gallbladder is justifiable. *Surgeon* 2003; 1: 233–235.
37. Grobmyer SR, Lieberman MD, Daly JM. Gallbladder cancer in the twentieth century: single institution's experience. *World J Surg* 2004; 28: 47–49.
38. Solaini L, Sharma A, Watt J et al. Predictive factors for incidental gallbladder dysplasia and carcinoma. *J Surg Res* 2014; 189: 17–21.
39. Sasatomi E, Tokunaga O, Miyazaki K. Precancerous conditions of gallbladder carcinoma: overview of histopathologic characteristics and molecular genetic findings. *J Hepatobiliary Pancreat Surg* 2000; 7: 556–567.
40. Hart J, Modan B, Shani M. Cholelithiasis in the aetiology of gallbladder neoplasms. *Lancet* 1971; 1: 1,151–1,153.
41. Sujata J, Rana S, Sabina K et al. Incidental gall bladder carcinoma in laparoscopic cholecystectomy: a report of 6 cases and a review of the literature. *J Clin Diagn Res* 2013; 7: 85–88.
42. Deng YL, Xiong XZ, Zhou Y et al. Selective histology of cholecystectomy specimens—is it justified? *J Surg Res* 2015; 193: 196–201.
43. Swank HA, Mulder IM, Hop WC et al. Routine histopathology for carcinoma in cholecystectomy specimens not evidence based: a systematic review. *Surg Endosc* 2013; 27: 4,439–4,448.
44. Bartlett DL, Fong Y, Fortner JG et al. Long-term results after resection for gallbladder cancer. Implications for staging and management. *Ann Surg* 1996;224: 639–646.
45. Choi SB, HJ H, Kim CY, et al. Incidental gallbladder cancer diagnosed following laparoscopic cholecystectomy. *W J Surg* 2009; 33(12):2657–2663.